Derm'Expert

INFANTILE HEMANGIOMA

5% of infants below the age of 1 year [2,5,6]




3 TYPES of Infantile Hemangioma (1)

SUPERFICIAL IH






MIXED/COMBINED IH





SUBCUTANEOUS/DEEP IH

DISEASE EVOLUTION (2)

3 PHASES




Pierre Fabre DERMATOLOGIE | SKIN EXPERTISE IN OUR DNA

INFANTILE HEMANGIOMA

SEQUELAE AFTER INVOLUTION OF UNTREATED HEMANGIOMAS [3-4]


Anatomic Distortion



Telangiectasias



Anetodermia/Redundant skin



Fibrofatty Tissue



Scar



1. Leauté-Labrèze C, Prey S, Ezzedine K. Infantile haemangioma: Part I. Pathophysiology, epidemiology, clinical features, life cycle and associated structural abnormalities. J Eur Acad Dermatol Venereol. 2011 Nov;25(11):1245-53.
2. Leauté-Labrèze C, Harper JI, Hoeger PH. Infantile haemangioma. The Lancet. 2017 Jul;390(10089):85-94.
3. Darrow D, Greene AK, Mancini AJ, Nopper A J. Diagnosis and Management of Infantile Hemangioma. PEDIATRICS. 2015 Oct;136(4):1060-1104.
4. Baselga E, Roe E, Coulie J, Munoz Fania Z, Boon LM, McCuaig C et al. Risk Factors for Degree and Type of Sequelae After Involution of Untreated Hemangiomas of Infancy. JAMA Dermatol. 2016 Nov;152(11):1239-1243.
5. Munden et al – Incidence and demographics of IH – BJD 2014, 170; 907-913
6. Volc-Platzer et al. – Poster ESPD 2014 – 4. Chen et al. Update on pathogenesis and therapy. Pediatrics. 2012

Pictures source: PFD internal images bank

FR TERRES DU SUD HQ-HEM-05-21-2100006 - © Istock

Pierre Fabre DERMATOLOGIE | SKIN EXPERTISE IN OUR DNA