


Instructions for use

Adapted from Instructions for Use approved by the U.S. Food and Drug Administration

1

Remove the items from the box


- Place your box of supplies on a clean flat work surface, such as a table.
- Remove the HEMANGEOL® bottle and oral dosing syringe from the box. **Do not shake the bottle before use.** Keep the box for storage.



2

Check the dose


- Remove the oral dosing syringe from the plastic bag.
- Safely throw the plastic bag away. The barrel of the syringe has markings in milliliters (mL).
- Look at the markings on the barrel of the oral dosing syringe and find the mL marking that matches the HEMANGEOL dose in mL prescribed by your doctor.



5

Remove the dose


- Use one hand to hold the oral dosing syringe in place. With your other hand, turn the bottle upside down.
- Pull back on the plunger until the top of the plunger lines up with the marking on the barrel of the syringe that matches the dose of HEMANGEOL prescribed by your doctor. Your child's dose may be different than the dose shown in this picture.



8

Give HEMANGEOL to your child

- Slowly squirt HEMANGEOL into your child's mouth after placing the oral dosing syringe against the inside of the cheek.
- Keep your child in an upright position for a few minutes right after giving a dose of HEMANGEOL.



- If needed, you can dilute the dose of HEMANGEOL in a small amount of milk or fruit juice and give it to your child in a baby's bottle. If your child spits up a dose or if you are not sure your child got all of the medicine, do not give another dose. Wait until the next scheduled dose.

3

Open the bottle

- Open the bottle of HEMANGEOL by pushing down on the plastic cap while turning the cap to the left.
- Write down on the box the date when you first open the bottle.



6

Check for air bubbles

- Check for air bubbles in the oral dosing syringe.
- If you see air bubbles, push up on the plunger towards the bottle just enough to remove any large air bubbles and then pull back to the measured dose.



9

Close the bottle


- Replace the plastic cap on the bottle by turning it clockwise (to the right).



4

Insert the syringe


- Place the bottle on your work surface. Use one hand to hold the bottle upright. Use your other hand to insert the tip of the oral dosing syringe into the syringe adapter at the top of the bottle. Push the plunger all the way down.
- Do not remove the syringe adapter. If the syringe adapter is missing talk to your pharmacist.



7

Remove the syringe

- Turn bottle upright again and place it on your work surface. Remove the oral dosing syringe from the bottle. Do not push the plunger in during this step.
- The syringe adapter should stay attached to the bottle.



10

Clean the syringe

- Clean the oral dosing syringe after each use by rinsing with clean tap water.
- Do not take apart the oral dosing syringe.
- Do not use any soap or alcohol based product to clean. Wipe the outside dry.
- Do not put the oral dosing syringe through a sterilizer or dishwasher.

